



INDICATORS OF PRIMARY HEALTH CARE

Definition of an indicator:

- Measurable variable used as a representation of an associated factor or quantity.
- Health indicators are summary measures that are designed to describe particular aspects of health or health system performance.

- A health indicator is a single measure that is reported on regularly and that provides relevant information about population health and/or health system performance and characteristics.



✓ Indicators related to the different primary health care activities:

I-Utilization rates

II- Indicators of quality of life

III- Mortality and morbidity indices.

I-Utilization rates

Indicators	PHC activity
I-Utilization rates	PHC services utilized.
1- Percentage of children below two years who are fully immunized with EPI vaccines.	<ul style="list-style-type: none">•EPI•Health education.•Under five health services.
2- Percentage of pregnant women attending antenatal clinics at least 5 times as per schedule. (3,4,5,8,9 th month)	<ul style="list-style-type: none">•MCH (antenatal care)•Health education.
3- Percentage of unattended home deliveries.	<ul style="list-style-type: none">•MCH (antenatal and intranatal care)•Health education.

4- Percentage of pregnant mothers fully immunized with TT.	<ul style="list-style-type: none"> •EPI •MCH (Antenatal care) • Health education.
5- Percentage of pregnant women receiving at least one postnatal home visit.	<ul style="list-style-type: none"> •MCH (Postnatal care, domiciliary service)
6- Percentage of high risk cases detected	<ul style="list-style-type: none"> •MCH (Antenatal)
7- Percentage of high risk cases referred to hospital (at least once)	<ul style="list-style-type: none"> •MCH (Antenatal, referral, equity)
8- Percentage of complicated deliveries	<ul style="list-style-type: none"> •MCH (Antenatal, postnatal, referral)
9- Percentage of receiving routine iron.	<ul style="list-style-type: none"> •Nutritional •MCH (Antenatal)

II- indicators of quality of life

Indicators	PHC activity
II- indicators of quality of life	Promotive, preventive services, community participation, intersectoral action
1. Percentage of infants with birth weight more than 2500gm.	<ul style="list-style-type: none">•Nutrition• MCH (antenatal)•Health education
2. Percentage of mothers breast-feeding infants for 1 year or more.	<ul style="list-style-type: none">•Nutrition• MCH•Health education

3. Percentage of mothers giving supplementary food at age of six month.

- Health education
- MCH
- Nutrition

4. Percentage of houses having round the year potable water supply.

- Sanitation
- Community participation
- Intersectoral coordination
- Health education
- Disease control

5. Percentage of houses having clean latrines.

- Sanitation
- Health education
- Diseases control.

6. Percentage of children receiving primary education.

- Intersectoral.

7. Percentage of rural houses having separate animal dwelling away from the house.

- Sanitation,
- Diseases control
- Community participation.

8. Life expectancy at birth

- PHC services (cumulative).

III. Mortality and morbidity indices

Indicators	PHC activity
III. Mortality and morbidity indices.	Equity, quality, diseases control
1. Infant mortality rate	<ul style="list-style-type: none">•MCH•EPI•Sanitation•CDC•Curative services
2. Maternal mortality rate	<ul style="list-style-type: none">•MCH•Nutrition•Referral•CDC

3. Perinatal mortality rate

- MCH (Antenatal, intranatal, postnatal)
- Nutrition
- Secondary care support

4. Neonatal mortality rate

- MCH (Intranatal, postnatal)
- EPI
- CDC
- Curative services

5. Incidence rate related to EPI diseases

- EPI
- CDC
- Sanitation
- MCH

6. 1-4 year mortality rate

- MCH
- CDC
- EPI
- Sanitation
- Health education
- Curative services

7. Diarrheal cases/deaths
below 5 years

- CDC
- Sanitation
- Health education
- Curative services



THANK YOU